








## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>5</sup> : C07D 209/14, A61K 31/40, C07D 401/06		A1	(11) International Publication Number: <b>WO 94/24105</b>
			(43) International Publication Date: 27 October 1994 (27.10.94)
(21) International Application Number: PCT/GB94/00760 (22) International Filing Date: 8 April 1994 (08.04.94) (30) Priority Data: 9307831.9      15 April 1993 (15.04.93)      GB 9316274.1      5 August 1993 (05.08.93)      GB (71) Applicant (for all designated States except US): MERCK SHARP & DOHME LIMITED [GB/GB]; Hertford Road, Hoddesdon, Hertfordshire EN11 9BU (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): KULAGOWSKI, Janusz, Jozef [GB/GB]; Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR (GB). LEESON, Paul, David [GB/GB]; Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR (GB). RIDGILL, Mark, Peter [GB/GB]; Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR (GB). (74) Agent: THOMPSON, John; Merck & Co., Inc., Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR (GB).			(81) Designated States: AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, HU, JP, KP, KR, KZ, LK, LU, LV, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  Published With international search report.

(54) Title: INDOLE DERIVATIVES AS DOPAMINE D<sub>4</sub> ANTAGONISTS

## (57) Abstract

Compound of formula (I), or a pharmaceutically acceptable salt thereof or a prodrug thereof wherein E represents -CH<sub>2</sub>- or -CH<sub>2</sub>CH<sub>2</sub>-; R represents hydrogen or C<sub>1-6</sub> alkyl; Q represents a moiety of formula Qa, Qb, Qc or Qd which are antagonists of dopamine receptor subtypes within the brain, having a selective affinity for the dopamine D<sub>4</sub> receptor subtype over other dopamine receptor subtypes, and are accordingly of benefit in the treatment and/or prevention of psychotic disorders such as schizophrenia whilst manifesting fewer side-effects than those associated with classical neuroleptic drugs.



**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

## INDOLE DERIVATIVES AS DOPAMINE D4 ANTAGONISTS

This invention relates to the use of a particular class of heteroaromatic compounds. More particularly, the invention is concerned with the use of substituted indole derivatives which are antagonists of dopamine receptor subtypes within the brain and are therefore of benefit in the treatment and/or prevention of psychotic disorders such as schizophrenia.

The "dopamine hypothesis" of schizophrenia predicts an increased activity of dopamine neurotransmission in the disease. The hypothesis is supported by early observations that drugs, such as amphetamine, with dopamine agonist or dopamine-releasing properties are capable of eliciting a psychosis indistinguishable from acute paranoid schizophrenia.

Schizophrenia is a disorder which is conventionally treated with drugs known as neuroleptics. In the majority of cases, the symptoms of schizophrenia can be treated successfully with so-called "classical" neuroleptic agents such as haloperidol. Classical neuroleptics generally are antagonists at dopamine D<sub>2</sub> receptors. The fact that classical neuroleptic drugs have an action on dopamine receptors in the brain thus lends credence to the "dopamine hypothesis" of schizophrenia.

Molecular biological techniques have revealed the existence of several subtypes of the dopamine receptor. The dopamine D<sub>1</sub> receptor subtype has been shown to occur in at least two discrete forms. Two forms of the D<sub>2</sub> receptor subtype, and at least one form of the D<sub>3</sub> receptor subtype, have also been discovered. More recently, the D<sub>4</sub> (Van Tol *et al.*, *Nature* (London), 1991, 350, 610) and D<sub>5</sub> (Sunahara *et al.*, *Nature* (London), 1991, 350, 614) receptor subtypes have been described.

- 2 -

Notwithstanding their beneficial antipsychotic effects, classical neuroleptic agents such as haloperidol are frequently responsible for eliciting acute extrapyramidal symptoms and neuroendocrine disturbances.

5 These side-effects, which clearly detract from the clinical desirability of classical neuroleptics, are believed to be attributable to D<sub>2</sub> receptor blockade in the striatal region of the brain. It is considered (Van Tol et al., supra) that compounds which can interact

10 selectively with the dopamine D<sub>4</sub> receptor subtype, whilst having a less-pronounced action at the D<sub>2</sub> subtype, might be free from, or at any rate less prone to, the side-effects associated with classical neuroleptics, whilst at the same time maintaining a beneficial level of

15 antipsychotic activity.

The compounds of use in the present invention, being antagonists of dopamine receptor subtypes within the brain, are accordingly of benefit in the treatment and/or prevention of psychotic disorders such as

20 schizophrenia. Moreover, the compounds of use in the invention have a selective affinity for the dopamine D<sub>4</sub> receptor subtype over other dopamine receptor subtypes, in particular the D<sub>2</sub> subtype, and can therefore be expected to manifest fewer side-effects than those

25 associated with classical neuroleptic drugs.

US Patent 3188313 describes inter alia certain 1-[(2-indolyl)-lower-alkyl]-4-substituted-piperazines, which are alleged therein to possess a panoply of depressant actions on the central and autonomic nervous

30 system, the cardiovascular system and the skeletal-muscular system, including blood pressure lowering, vomiting incidence decreasing, sleeping time potentiating, tranquilizing and skeletal muscle relaxing activities. They are consequently alleged to be useful

35 as hypotensive agents, antinauseants, antipyretics, sedatives, tranquilizers and skeletal muscle relaxants.


- 3 -

In addition, NL-A-6511642 describes a family of 2-[piperidinoalkyl]-1H-indole derivatives, which are stated therein to have analgesic and muscle relaxant activity.

5           There is, however, no suggestion in US-3188313 or NL-A-6511642 that the compounds described therein would be of any benefit in the treatment and/or prevention of psychotic disorders such as schizophrenia, still less that in doing so they might be expected to  
10       manifest fewer side-effects than those exhibited by classical neuroleptic agents.

The present invention accordingly provides the use of a compound of formula I, or a pharmaceutically acceptable salt thereof or a prodrug thereof:

15



( I )

wherein

E represents -CH<sub>2</sub>- or -CH<sub>2</sub>CH<sub>2</sub>-;





R represents hydrogen or C<sub>1-6</sub> alkyl;

Q represents a moiety of formula Qa, Qb, Qc or

30       Qd:

35

- 4 -



in which the broken line represents an optional chemical bond;

20  $R^1$  represents hydrogen, or an optionally substituted  $C_{1-6}$  alkyl,  $C_{1-6}$  alkoxy,  $C_{2-6}$  alkenyl,  $C_{2-6}$  alkynyl, aryl, aryl( $C_{1-6}$ )alkyl, aryl( $C_{1-6}$ )alkoxy, aryl( $C_{2-6}$ )alkenyl, aryl( $C_{2-6}$ )alkynyl,  $C_{3-7}$  heterocycloalkyl( $C_{1-6}$ )alkyl, heteroaryl, heteroaryl( $C_{1-6}$ )alkyl, heteroaryl( $C_{2-6}$ )alkenyl or heteroaryl( $C_{2-6}$ )alkynyl group;

25  $R^2$  represents an optionally substituted  $C_{1-6}$  alkyl,  $C_{1-6}$  alkoxy,  $C_{2-6}$  alkenyl,  $C_{2-6}$  alkynyl, aryl, aryl( $C_{1-6}$ )alkyl, aryl( $C_{1-6}$ )alkoxy, aryl( $C_{2-6}$ )alkenyl, aryl( $C_{2-6}$ )alkynyl,  $C_{3-7}$  heterocycloalkyl( $C_{1-6}$ )alkyl, heteroaryl, heteroaryl( $C_{1-6}$ )alkyl, heteroaryl( $C_{2-6}$ )alkenyl or heteroaryl( $C_{2-6}$ )alkynyl group;

30  $R^3$ ,  $R^4$  and  $R^5$  independently represent hydrogen, hydrocarbon, a heterocyclic group, halogen, cyano, trifluoromethyl, nitro,  $-OR^a$ ,  $-SR^a$ ,  $-SOR^a$ ,  $-SO_2R^a$ ,  
35

- 5 -

$-\text{SO}_2\text{NR}^a\text{R}^b$ ,  $-\text{NR}^a\text{R}^b$ ,  $-\text{NR}^a\text{COR}^b$ ,  $-\text{NR}^a\text{CO}_2\text{R}^b$ ,  $-\text{COR}^a$ ,  $-\text{CO}_2\text{R}^a$  or  $-\text{CONR}^a\text{R}^b$ ;

Z represents  $-\text{CH}_2-$  or  $-\text{CH}_2\text{CH}_2-$ ;

5  $\text{R}^6$  represents hydrogen,  $\text{C}_{1-6}$  alkyl,  $\text{C}_{1-6}$  alkoxy, aryl, aryl( $\text{C}_{1-6}$ )alkyl or halogen; and

$\text{R}^a$  and  $\text{R}^b$  independently represent hydrogen, hydrocarbon or a heterocyclic group; for the manufacture of a medicament for the treatment and/or prevention of psychotic disorders such as  
10 schizophrenia.

For use in medicine, the salts of the compounds of formula I will be pharmaceutically acceptable salts. Other salts may, however, be useful in the preparation of the compounds of use in the invention or of their  
15 pharmaceutically acceptable salts. Suitable pharmaceutically acceptable salts of the compounds of use in this invention include acid addition salts which may, for example, be formed by mixing a solution of the compound of use in the invention with a solution of a  
20 pharmaceutically acceptable acid such as hydrochloric acid, sulphuric acid, fumaric acid, maleic acid, succinic acid, acetic acid, benzoic acid, oxalic acid, citric acid, tartaric acid, carbonic acid or phosphoric acid. Furthermore, where the compounds of use in the invention  
25 carry an acidic moiety, suitable pharmaceutically acceptable salts thereof may include alkali metal salts, e.g. sodium or potassium salts; alkaline earth metal salts, e.g. calcium or magnesium salts; and salts formed with suitable organic ligands, e.g. quaternary ammonium  
30 salts.

The term "hydrocarbon" as used herein includes straight-chained, branched and cyclic groups containing up to 18 carbon atoms, suitably up to 15 carbon atoms, and conveniently up to 12 carbon atoms. Suitable  
35 hydrocarbon groups include  $\text{C}_{1-6}$  alkyl,  $\text{C}_{2-6}$  alkenyl,  $\text{C}_{2-6}$  alkynyl,  $\text{C}_{3-7}$  cycloalkyl,  $\text{C}_{3-7}$  cycloalkyl( $\text{C}_{1-6}$ )alkyl,

- 6 -

aryl, aryl(C<sub>1-6</sub>)alkyl, aryl(C<sub>2-6</sub>)alkenyl and aryl(C<sub>2-6</sub>)alkynyl.

5 The expression "a heterocyclic group" as used herein includes cyclic groups containing up to 18 carbon atoms and at least one heteroatom preferably selected from oxygen, nitrogen and sulphur. The heterocyclic group suitably contains up to 15 carbon atoms and conveniently up to 12 carbon atoms, and is preferably linked through carbon. Examples of suitable heterocyclic  
10 groups include C<sub>3-7</sub> heterocycloalkyl, C<sub>3-7</sub> heterocycloalkyl(C<sub>1-6</sub>)alkyl, heteroaryl, heteroaryl(C<sub>1-6</sub>)alkyl, heteroaryl(C<sub>2-6</sub>)alkenyl and heteroaryl(C<sub>2-6</sub>)alkynyl groups.

Suitable alkyl groups within the scope of the  
15 term "hydrocarbon" and within the definition of the substituents R, R<sup>1</sup>, R<sup>2</sup> and R<sup>6</sup> include straight-chained and branched alkyl groups containing from 1 to 6 carbon atoms. Typical examples include methyl and ethyl groups, and straight-chained or branched propyl and butyl groups.  
20 Particular alkyl groups are methyl, ethyl, n-propyl, isopropyl and t-butyl.

Suitable alkenyl groups within the scope of the term "hydrocarbon" and within the definition of the substituents R<sup>1</sup> and R<sup>2</sup> include straight-chained and  
25 branched alkenyl groups containing from 2 to 6 carbon atoms. Typical examples include vinyl and allyl groups.

Suitable alkynyl groups within the scope of the term "hydrocarbon" and within the definition of the substituents R<sup>1</sup> and R<sup>2</sup> include straight-chained and  
30 branched alkynyl groups containing from 2 to 6 carbon atoms. Typical examples include ethynyl and propargyl groups.

Suitable cycloalkyl groups include groups containing from 3 to 7 carbon atoms. Particular  
35 cycloalkyl groups are cyclopropyl and cyclohexyl.



- 7 -

Particular aryl groups within the scope of the term "hydrocarbon" and within the definition of the substituents  $R^1$ ,  $R^2$  and  $R^6$  include phenyl and naphthyl.

Particular aryl( $C_{1-6}$ )alkyl groups within the scope of the term "hydrocarbon" and within the definition of the substituents  $R^1$ ,  $R^2$  and  $R^6$  include benzyl, naphthylmethyl, phenethyl and phenylpropyl.

Suitable heterocycloalkyl groups include azetidiny, pyrrolidyl, piperidyl, piperazinyl, morpholinyl and tetrahydrofuryl groups.

A particular  $C_{3-7}$  heterocycloalkyl( $C_{1-6}$ )alkyl group within the scope of the expression "a heterocyclic group" and within the definition of the substituents  $R^1$  and  $R^2$  is tetrahydrofurylethyl.

Suitable heteroaryl groups within the scope of the expression "a heterocyclic group" and within the definition of the substituents  $R^1$  and  $R^2$  include pyridyl, quinolyl, isoquinolyl, pyridazinyl, pyrimidinyl, pyrazinyl, pyranyl, furyl, benzofuryl, dibenzofuryl, thienyl, benzthienyl, indolyl, indazolyl, imidazolyl, benzimidazolyl, oxadiazolyl and thiadiazolyl groups.

Particular heteroaryl( $C_{1-6}$ )alkyl groups within the scope of the expression "a heterocyclic group" and within the definition of the substituents  $R^1$  and  $R^2$  include thienylmethyl, pyridylmethyl, pyrimidinylmethyl and pyrazinylmethyl.

The hydrocarbon and heterocyclic groups, as well as the substituents  $R^1$  and  $R^2$ , may in turn be optionally substituted by one or more groups selected from  $C_{1-6}$  alkyl, adamantyl, phenyl, aryl( $C_{1-6}$ )alkyl, halogen,  $C_{1-6}$  haloalkyl,  $C_{1-6}$  aminoalkyl, trifluoromethyl, hydroxy,  $C_{1-6}$  alkoxy, aryloxy, keto,  $C_{1-3}$  alkylenedioxy, nitro, cyano, carboxy,  $C_{2-6}$  alkoxycarbonyl,  $C_{2-6}$  alkoxycarbonyl( $C_{1-6}$ )alkyl,  $C_{2-6}$  alkylcarbonyloxy, arylcarbonyloxy,  $C_{2-6}$  alkylcarbonyl, arylcarbonyl,  $C_{1-6}$  alkylthio,  $C_{1-6}$  alkylsulphinyl,  $C_{1-6}$

- 8 -

alkylsulphonyl, arylsulphonyl, trifluoromethane-sulphonyloxy,  $-NR^V R^W$ ,  $-NR^V COR^W$ ,  $-NR^V CO_2 R^W$ ,  $-NR^V SO_2 R^W$ ,  $-CH_2 NR^V SO_2 R^W$ ,  $-NHCONR^V R^W$ ,  $-PO(OR^V)(OR^W)$ ,  $-CONR^V R^W$ ,  $-SO_2 NR^V R^W$  and  $-CH_2 SO_2 NR^V R^W$ , in which  $R^V$  and  $R^W$  independently represent hydrogen,  $C_{1-6}$  alkyl, aryl or aryl( $C_{1-6}$ )alkyl.

The term "halogen" as used herein includes fluorine, chlorine, bromine and iodine, especially chlorine.

The present invention includes within its scope the use of prodrugs of the compounds of formula I above. In general, such prodrugs will be functional derivatives of the compounds of formula I which are readily convertible in vivo into the required compound of formula I. Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in "Design of Prodrugs", ed. H. Bundgaard, Elsevier, 1985.

Where the compounds of use in the invention have at least one asymmetric centre, they may accordingly exist as enantiomers. Where the compounds of use in the invention possess two or more asymmetric centres, they may additionally exist as diastereoisomers. It is to be understood that the use of all such isomers and mixtures thereof is encompassed within the scope of the present invention.

Suitably, the substituent R represents hydrogen or methyl, especially hydrogen.

Suitably, the substituent  $R^1$  represents hydrogen.

Suitable values for the substituent  $R^2$  include  $C_{1-6}$  alkyl, phenyl, halophenyl,  $C_{1-6}$  alkylphenyl,  $C_{1-6}$  alkoxyphenyl, nitrophenyl, benzyl, halobenzyl, phenethyl, phenylpropyl and benzyloxy. Particular values of  $R^2$  include methyl, ethyl, n-propyl, isopropyl, phenyl, chlorophenyl, ethylphenyl, methoxyphenyl, nitrophenyl,


- 9 -

benzyl, chlorobenzyl, phenethyl, phenylpropyl and benzyloxy.

Suitable values for the substituents  $R^3$ ,  $R^4$  and  $R^5$  include hydrogen, halogen, cyano, nitro, trifluoromethyl, amino,  $C_{1-6}$  alkylamino, di( $C_{1-6}$ )alkylamino,  $C_{1-6}$  alkyl,  $C_{1-6}$  alkoxy, aryl( $C_{1-6}$ )alkoxy and  $C_{2-6}$  alkylcarbonyl. Particular values include hydrogen, fluoro, chloro, methyl, methoxy and benzyloxy.

Particular values of  $R^6$  include hydrogen, phenyl, chloro and bromo.

A particular sub-class of compounds of use in the invention is represented by the compounds of formula IIA, and pharmaceutically acceptable salts thereof and prodrugs thereof:



(IIA)

wherein

$E$  is as defined with reference to formula I above;

$n$  is zero, 1, 2 or 3;

$W$  is CH or nitrogen; and


$R^{13}$  and  $R^{17}$  independently represent hydrogen, halogen, cyano, nitro, trifluoromethyl, amino,  $C_{1-6}$  alkylamino, di( $C_{1-6}$ )alkylamino,  $C_{1-6}$  alkyl,  $C_{1-6}$  alkoxy, aryl( $C_{1-6}$ )alkoxy or  $C_{2-6}$  alkylcarbonyl.

Particular values of  $R^{13}$  include hydrogen, fluoro, chloro, methyl, ethyl, methoxy and benzyloxy.

- 10 -

Particular values of  $R^{17}$  include hydrogen, chloro, methoxy and nitro.

Another sub-class of compounds of use in the invention is represented by the compounds of formula IIB, and pharmaceutically acceptable salts thereof and prodrugs thereof:




(IIB)

wherein

E is as defined with reference to formula I above; and

n,  $R^{13}$  and  $R^{17}$  are as defined with reference to formula IIA above.

A further sub-class of compounds of use in the invention is represented by the compounds of formula IIC, and pharmaceutically acceptable salts thereof and prodrugs thereof:



(IIC)

wherein

35

m is 1, 2 or 3;


-X-Y- represents  $-CH_2CH-$  or  $-CH=C-$ ;

- 11 -

E is as defined with reference to formula I above; and

$R^{13}$  and  $R^{17}$  are as defined with reference to formula IIA above.

5 A still further sub-class of compounds of use in the invention is represented by the compounds of formula IID, and pharmaceutically acceptable salts thereof and prodrugs thereof:



(IID)

wherein

20 E is as defined with reference to formula I above;

Z represents  $-CH_2-$  or  $-CH_2CH_2-$ ;

$R^{13}$  is as defined with reference to formula IIA above; and


25  $R^{16}$  represents hydrogen,  $C_{1-6}$  alkyl,  $C_{1-6}$  alkoxy, aryl, aryl( $C_{1-6}$ )alkyl or halogen.

Particular values of  $R^{16}$  include hydrogen, phenyl, chloro and bromo, especially hydrogen.

30 A yet further sub-class of compounds of use in the invention is represented by the compounds of formula IIE, and pharmaceutically acceptable salts thereof and prodrugs thereof:

35

- 12 -



(IIE)

10 wherein

E is as defined with reference to formula I above;

R<sup>13</sup> and R<sup>17</sup> are as defined with reference to formula IIA above; and

15 m is 1, 2 or 3.

Specific compounds of use in the present invention include:

2-(4-phenylpiperazin-1-ylmethyl)-1H-indole;

2-[4-(4-chlorophenyl)piperazin-1-ylmethyl]-1H-indole;

20 and pharmaceutically acceptable salts thereof and prodrugs thereof.

Certain compounds falling within the definition of formula I above are novel. Particular sub-classes of novel compounds in accordance with the present invention  
25 comprise the compounds of formulae IIB, IIC, IID and IIE as defined above, and salts and prodrugs thereof. The invention further provides a novel compound selected from the following:

2-(4-benzylpiperazin-1-ylmethyl)-1H-indole;

30 2-[2-(4-benzylpiperazin-1-yl)ethyl]-1H-indole;

2-[4-(4-methoxyphenyl)piperazin-1-ylmethyl]-1H-indole;

2-(4-phenyl-1,2,3,6-tetrahydropyrid-1-ylmethyl)-1H-indole;

2-[4-(2-phenylethyl)piperidin-1-ylmethyl]-1H-indole;

35 2-(4-benzylloxypiperidin-1-ylmethyl)-1H-indole;

- 13 -

2-[4-(2-phenylethyl)-1,2,3,6-tetrahydropyrid-1-ylmethyl]-1H-indole;

2-[4-(3-phenylpropyl)-1,2,3,6-tetrahydropyrid-1-ylmethyl]-1H-indole;

5 2-[4-(3-phenylpropylidene)piperidin-1-ylmethyl]-1H-indole;

2-(1,2,3,4-tetrahydroisoquinolin-2-ylmethyl)-1H-indole;

2-(1,2-dihydroisoindol-2-ylmethyl)-1H-indole;

and salts and prodrugs thereof.

10           The invention also provides pharmaceutical compositions comprising one or more of the novel compounds according to the invention in association with a pharmaceutically acceptable carrier. Preferably these compositions are in unit dosage forms such as tablets,  
15 pills, capsules, powders, granules, sterile parenteral solutions or suspensions, metered aerosol or liquid sprays, drops, ampoules, auto-injector devices or suppositories; for oral, parenteral, intranasal, sublingual or rectal administration, or for  
20 administration by inhalation or insufflation. Alternatively, the compositions may be presented in a form suitable for once-weekly or once-monthly administration; for example, an insoluble salt of the active compound, such as the decanoate salt, may be  
25 adapted to provide a depot preparation for intramuscular injection. For preparing solid compositions such as tablets, the principal active ingredient is mixed with a pharmaceutical carrier, e.g. conventional tableting ingredients such as corn starch, lactose, sucrose,  
30 sorbitol, talc, stearic acid, magnesium stearate, dicalcium phosphate or gums, and other pharmaceutical diluents, e.g. water, to form a solid preformulation composition containing a homogeneous mixture of a compound of the present invention, or a pharmaceutically acceptable salt thereof. When referring to these  
35 preformulation compositions as homogeneous, it is meant

- 14 -

that the active ingredient is dispersed evenly throughout the composition so that the composition may be readily subdivided into equally effective unit dosage forms such as tablets, pills and capsules. This solid

5 preformulation composition is then subdivided into unit dosage forms of the type described above containing from 0.1 to about 500 mg of the active ingredient of the present invention. The tablets or pills of the novel

10 composition can be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope

15 over the former. The two components can be separated by an enteric layer which serves to resist disintegration in the stomach and permits the inner component to pass intact into the duodenum or to be delayed in release. A variety of materials can be used for such enteric layers or coatings, such materials including a number of

20 polymeric acids and mixtures of polymeric acids with such materials as shellac, cetyl alcohol and cellulose acetate.

The liquid forms in which the novel compositions of the present invention may be incorporated

25 for administration orally or by injection include aqueous solutions, suitably flavoured syrups, aqueous or oil suspensions, and flavoured emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil or peanut oil, as well as elixirs and similar pharmaceutical

30 vehicles. Suitable dispersing or suspending agents for aqueous suspensions include synthetic and natural gums such as tragacanth, acacia, alginate, dextran, sodium carboxymethylcellulose, methylcellulose, polyvinyl-pyrrolidone or gelatin.


35 In the treatment of schizophrenia, a suitable dosage level is about 0.01 to 250 mg/kg per day,




- 15 -

preferably about 0.05 to 100 mg/kg per day, and especially about 0.05 to 5 mg/kg per day. The compounds may be administered on a regimen of 1 to 4 times per day.

The compounds of formula I above, including the novel compounds according to the present invention, may be prepared by a process which comprises reacting a compound of formula III with a compound of formula IV:



(III)



(IV)


wherein E, R, R<sup>3</sup>, R<sup>4</sup> and R<sup>5</sup> are as defined above, Q<sup>1</sup> represents the residue of a moiety of formula Qa to Qd as defined above, and L represents a suitable leaving group.

The leaving group L is suitably a halogen atom, e.g. chlorine or bromine.

The reaction is conveniently carried out by stirring the reactants under basic conditions in a suitable solvent, for example potassium carbonate in N,N-dimethylformamide, or triethylamine in tetrahydrofuran or acetonitrile.

In an alternative procedure, the compounds of formula I above, including the novel compounds according to the present invention, may be prepared by a process which comprises reducing a compound of formula V:

- 16 -




(V)

10 wherein  $Q^1$ , R,  $R^3$ ,  $R^4$  and  $R^5$  are as defined above; and  $E^1$  represents a bond or a methylene group.

The reaction is conveniently carried out by treating the compound V with a reducing agent such as lithium aluminium hydride in an appropriate solvent, e.g.  
 15 tetrahydrofuran.

The intermediates of formula V above may suitably be prepared by reacting a compound of formula IV as defined above with a compound of formula VI:



(VI)

wherein R,  $R^3$ ,  $R^4$ ,  $R^5$  and  $E^1$  are as defined above; and G represents a reactive carboxylate moiety.


30 Suitable values for the reactive carboxylate moiety G include esters, for example  $C_{1-4}$  alkyl esters; acid anhydrides, for example mixed anhydrides with  $C_{1-4}$  alkanolic acids; acid halides, for example acid chlorides; and acylimidazoles.

35 By way of example, the intermediates of formula VI above wherein G is an acid chloride moiety may be

- 17 -

prepared by treating the corresponding carboxylic acid derivative with thionyl chloride in toluene. Similarly, the intermediates of formula VI wherein G is an acylimidazole moiety may be prepared by treating the corresponding carboxylic acid derivative with 1,1'-carbonyldiimidazole. Alternatively, the reactive carboxylate moiety G may be obtained by treating the corresponding compound wherein G is carboxy with 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride and 1-hydroxybenzotriazole hydrate, optionally in the presence of triethylamine; the resulting activated carboxylate intermediate may then suitably be reacted in situ with the required compound of formula IV.

In a further procedure, the compounds of formula I above, including the novel compounds according to the invention, may be prepared by a process which comprises the ring closure of a compound of formula VII:



(VII)


wherein  $R$ ,  $R^3$ ,  $R^4$ ,  $R^5$ ,  $E$  and  $Q^1$  are as defined above; and  $T^-$  represents a suitable counterion, e.g. a halide ion such as bromide.

The ring closure reaction is conveniently carried out under conditions analogous to those described in Chem. Ber., 1986, 119, 2069-2074. This requires the presence of a strong base, typically potassium *t*-butoxide; and the reaction is suitably effected in an

- 18 -

inert organic solvent at an elevated temperature, for example toluene at reflux.


The intermediates of formula VII above may suitably be prepared by reacting a compound of formula IV  
5 as defined above with a compound of formula VIII:



(VIII)

15 wherein R, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, E, L and T are as defined above.

The intermediates of formula VIII may in turn be conveniently prepared by reacting a compound of formula IX:



(IX)

with an acylating agent of formula G-E-L; wherein R, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, E, L, G and T are as defined above.

30 Where they are not commercially available, the starting materials of formula III, IV, VI, IX and G-E-L may be prepared by procedures analogous to those described in the accompanying Examples, or by standard methods well known from the art.

35 It will be appreciated that any compound of formula I initially obtained from any of the above

- 19 -

processes may, where appropriate, subsequently be elaborated into a further desired compound of formula I using techniques known from the art. For example, a compound of formula I wherein R is hydrogen initially  
5 obtained may be converted into a compound of formula I wherein R represents C<sub>1-6</sub> alkyl by standard alkylation techniques, such as by treatment with an alkyl iodide, e.g. methyl iodide, typically under basic conditions, e.g. sodium hydride in dimethylformamide, or  
10 triethylamine in acetonitrile.

Where the above-described processes for the preparation of the compounds of use in the invention give rise to mixtures of stereoisomers, these isomers may be separated by conventional techniques such as preparative  
15 chromatography. The compounds may be prepared in racemic form, or individual enantiomers may be prepared either by enantiospecific synthesis or by resolution. The compounds may, for example, be resolved into their component enantiomers by standard techniques such as  
20 preparative HPLC, or the formation of diastereomeric pairs by salt formation with an optically active acid, such as (-)-di-p-toluoyl-d-tartaric acid and/or (+)-di-p-toluoyl-l-tartaric acid, followed by fractional crystallization and regeneration of the free base. The  
25 compounds may also be resolved by formation of diastereomeric esters or amides, followed by chromatographic separation and removal of the chiral auxiliary.

During any of the above synthetic sequences it  
30 may be necessary and/or desirable to protect sensitive or reactive groups on any of the molecules concerned. This may be achieved by means of conventional protecting groups, such as those described in Protective Groups in Organic Chemistry, ed. J.F.W. McOmie, Plenum Press, 1973; and T.W. Greene & P.G.M. Wuts, Protective Groups in Organic Synthesis, John Wiley & Sons, 1991. The  
35

- 20 -

protecting groups may be removed at a convenient subsequent stage using methods known from the art.

The following Examples illustrate the preparation of compounds according to the invention.

5           The compounds useful in this invention potentially inhibit [ $^3\text{H}$ ]-spiperone binding to human dopamine  $\text{D}_4$  receptor subtypes expressed in clonal cell lines.

#### [ $^3\text{H}$ ]-Spiperone Binding Studies

10           Clonal cell lines expressing the human dopamine  $\text{D}_4$  receptor subtype were harvested in PBS and then lysed in 10 mM Tris-HCl pH 7.4 buffer containing 5 mM  $\text{MgSO}_4$  for 20 min on ice. Membranes were centrifuged at 50,000g for 15 min at 4°C and the resulting pellets resuspended in assay buffer (50 mM Tris-HCl pH 7.4 containing 5 mM EDTA, 1.5 mM  $\text{CaCl}_2$ , 5 mM  $\text{MgCl}_2$ , 5 mM KCl, 120 mM NaCl, and 0.1% ascorbic acid) at 20 mg/ml wet weight. Incubations were carried out for 60 min at room temperature (22°C) in the presence of 0.05-2 nM [ $^3\text{H}$ ]-spiperone or 0.2 nM for displacement studies and were initiated by addition of 20-100  $\mu\text{g}$  protein in a final assay volume of 0.5 ml. The incubation was terminated by rapid filtration over GF/B filters presoaked in 0.3% PEI and washed with 10 ml ice-cold 50 mM Tris-HCl, pH 7.4. Specific binding was determined by 10  $\mu\text{M}$  apomorphine and radioactivity determined by counting in a LKB beta counter. Binding parameters were determined by non-linear least squares regression analysis, from which the inhibition constant  $\text{K}_i$  could be calculated for each test compound.

30           The compounds of the accompanying Examples were tested in the above assay, and all were found to possess a  $\text{K}_i$  value for displacement of [ $^3\text{H}$ ]-spiperone from the human dopamine  $\text{D}_4$  receptor subtype of below 1.5  $\mu\text{M}$ .

35

- 21 -

EXAMPLE 12-(2-[4-Phenylpiperazin-1-yl]ethyl)indole

Step 1: 2-(3-Chloropropionylamino)benzyltriphenylphosphonium bromide

5

3-Chloropropionyl chloride (3.0ml, 32mmol) and piperidine (3.5ml, 43.5mmol) were added to a suspension of 2-aminobenzyltriphenylphosphonium bromide (13.0g, 29mmol) (synthesised using the procedure of Capuano *et al*, Chem. Ber., 10 1986, 119, 2069-2074) in dichloromethane (60ml) under a nitrogen atmosphere. The mixture was heated at reflux for 30 min. and then cooled to room temperature. Dichloromethane (60ml) and hydrochloric acid (1N, 50ml) were added and the precipitated solid was collected by filtration, washed with 15 dichloromethane and dried *in vacuo* to give 2-(3-chloropropionylamino)benzyltriphenylphosphonium bromide (7.5g, 48%). The filtrate was diluted with dichloromethane (300ml) and washed with sodium carbonate (satd., 150ml). Evaporation of the dichloromethane gave a residue which was 20 triturated with ether (200ml) and the white solid collected to afford a second crop of product (5.0g, 32%).

Step 2: 2-(3-[4-phenylpiperazin-1-yl]propionylamino)benzyltriphenylphosphonium bromide

25

A mixture of 2-(3-chloropropionylamino)benzyltriphenylphosphonium bromide (2.70g, 5mmol) and phenylpiperazine (1.70g, 10mmol) in acetonitrile (85ml) was heated at reflux for 2.5h. The mixture was cooled to room temperature and filtered and the filtrate evaporated *in vacuo* to 30 give 2-(3-[4-phenylpiperazin-1-yl]propionylamino)benzyltriphenylphosphonium bromide as a white solid (3.1g, 93%).

- 22 -

Step 3: 2-(2-[4-Phenylpiperazin-1-yl]ethyl)indole

The phosphonium salt (3.1g, 4.6mmol) was suspended in toluene (150ml) and heated at reflux under Dean-Stark conditions for 90 min. Freshly dried potassium *tert*-butoxide (716mg, 5.5mmol) was added and the mixture was heated at reflux for 30 min. then filtered hot. The cooled filtrate was concentrated *in vacuo* to about 20ml. The residue was dissolved in dichloromethane (100ml) and extracted with hydrochloric acid (1M, 2 x 50ml). The dichloromethane extract was evaporated *in vacuo* and the residue purified by flash chromatography using ethyl acetate as eluant. Recrystallisation from methanol gave the *title compound* as a white solid (18mg, 1.3%), mp 184-186°C; (Found: C, 78.59; H, 7.56; N, 13.50. C<sub>20</sub>H<sub>23</sub>N<sub>3</sub> requires C, 78.65; H, 7.59; N, 13.76%);  $\delta_H$  (DMSO-d<sub>6</sub>) 2.59-2.61 (4H, m, 2 x piperazinyl CH<sub>2</sub>), 2.60 (2H, t, J 4.9Hz, CH<sub>2</sub>CH<sub>2</sub>N), 2.69-2.73 (2H, m, CH<sub>2</sub>CH<sub>2</sub>N), 3.14 (4H, t, J 4.9Hz, 2 x piperazinyl CH<sub>2</sub>), 6.17 (1H, s, 3-H), 6.76 (1H, t, J 7.2Hz, ArH), 6.78-7.00 (4H, m, ArH), 7.18-7.22 (2H, m, ArH), 7.27 (1H, d, J 8.0Hz, 4-H), 7.39 (1H, d, J 7.6Hz, 7-H), and 10.84 (1H, br s, NH); m/z (CI<sup>+</sup>, NH<sub>3</sub>) 306 (M+1)<sup>+</sup>.

EXAMPLE 22-(2-[4-Benzylpiperazin-1-yl]ethyl)indoleStep 1: Indole-2-acetic acid

A solution of lithium hydroxide (346mg, 8.25mmol) in water (2ml) was added to methyl indole-2-acetate (520mg, 2.75mmol) (prepared using the method of Capuano *et al*, Chem. Ber., 1986, 119, 2069) in THF (2ml) and methanol (6ml) and the mixture stirred for 1h. The mixture was concentrated *in vacuo*, diluted with water (20ml) and extracted with diethyl ether (2 x 10ml). The aqueous layer was carefully acidified to pH 2 with



- 23 -

1M HCl and extracted into dichloromethane (3 x 20ml). The extracts were dried ( $\text{MgSO}_4$ ) and evaporated *in vacuo* to give indole-2-acetic acid (350mg, 72%) as a pink solid.

5           Step 2: 1-benzyl-4-(2-indolylacetamido)piperazine

1-Hydroxybenzotriazole (324mg, 2.4mmol) and triethylamine (446 $\mu$ l, 3.2mmol) were added to a solution of indole-2-acetic acid (350mg, 2mmol) and 1-benzylpiperazine (352mg, 2mmol) in THF (25ml), followed after 5 min by 1-(3-dimethylaminopropyl)-3-ethyl carbodiimide hydrochloride (576mg, 3mmol) and the mixture stirred for 15 h. The mixture was concentrated *in vacuo* and the residue partitioned between citric acid (1M, 30ml) and ethyl acetate (25ml), the phases were separated and the aqueous layer extracted with ethyl acetate (2 x 25ml). The combined organic layers were washed with sodium bicarbonate (satd., 2 x 10ml) and brine (satd., 20ml), dried ( $\text{Na}_2\text{SO}_4$ ) and evaporated *in vacuo* to give 1-benzyl-4-(2-indolylacetamido)piperazine as a pink solid (200mg, 30%), m/z (CI<sup>+</sup>,  $\text{NH}_3$ ) 334 (M+1)<sup>+</sup>.

          Step 3: 2-(2-[4-Benzylpiperazin-1-yl]ethyl)indole

Lithium aluminium hydride in ether (1M, 600 $\mu$ l, 0.6mmol) was carefully added to a solution of 1-benzyl-4-(2-indolylacetamido)piperazine (200mg, 0.6mmol) in THF (10ml) and the mixture was stirred at reflux for 1 h. The reaction was cooled in ice and successively treated with water (0.5ml), sodium hydroxide (4M, 0.5ml) and water (1.5ml). The mixture was filtered and the filtrate concentrated *in vacuo*. The residue was purified by column chromatography on silica (90:8:1  $\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{NH}_3$ ), the resulting gum dissolved in methanol and ether and treated with oxalic acid to give a brown solid (32mg, 11%). M.p. 215-216°C (MeOH); (Found: C, 59.60; H, 5.99;

- 24 -

5 N, 7.92.  $C_{21}H_{25}N_3 \cdot 2(CO_2H)_2 \cdot 0.3H_2O$  requires C, 59.47; H, 5.91; N, 8.32%;  $\delta_H$  (DMSO- $d_6$ ) 2.78 (3H, br s, piperazinyl H), 3.06 (7H, m), 3.76 (2H, s,  $CH_2$ ), 6.22 (1H, s, 3-H), 6.93 (1H, t, J 7.3Hz, ArH), 7.01 (1H, m, ArH), 7.28-7.43 (7H, m, ArH), and 11.01 (1H, br s, NH);  $m/z$  ( $CI^+$ ,  $NH_3$ ) 320 ( $M+1$ ) $^+$ .

### EXAMPLE 3

#### 2-(4-Benzylpiperazin-1-ylmethyl)indole

10 Step 1: 2-(4-benzylpiperazin-1-yl)carbonyl indole hydrochloride

Thionyl chloride (1.62ml, 22.3mmol) was added to a solution of indole-2-carboxylic acid (3.0g, 18.6mmol) in toluene (40ml), the mixture heated at reflux for 3 h and then cooled to room temperature. 1-Benzylpiperazine (3.7ml, 20.4mmol) was added and the resultant precipitate collected by filtration, washed with toluene, and dried *in vacuo* to give 2-(4-benzylpiperazin-1-yl) carbonyl indole hydrochloride (6.5g, 98%). M.p. >210°C (dec).

20

#### Step 2: 2-(4-Benzylpiperazin-1-ylmethyl)indole

Lithium aluminium hydride in THF (1M, 11.25mmol) was slowly added to a suspension of 2-(4-benzylpiperazin-1-yl)carbonyl indole hydrochloride (2.0g, 5.6mmol) in THF (40ml) under a nitrogen atmosphere and the resultant solution heated at reflux for 90 min. The mixture was cooled to room temperature and treated sequentially with water (0.5ml), sodium hydroxide (4M, 0.5ml) and water (1.5ml). The mixture was filtered and the filtrate evaporated *in vacuo*. The residue was recrystallised from toluene to afford the *title compound* as a white solid (428mg, 25%), m.p. 157-159°C; (Found: C, 78.44; H,

25  
30

- 25 -

7.63; N, 13.66.  $C_{20}H_{23}N_3$  requires C, 78.65; H, 7.59; N, 13.76%;  $\delta_H$  (DMSO- $d_6$ ) 2.40 (8H, br s, 4 x piperazinyl  $CH_2$ ), 3.45 (2H, s,  $CH_2$ ), 3.59 (2H, s,  $CH_2$ ), 6.23 (1H, s, 3-H), 6.90 (1H, m, ArH), 6.95 (1H, m, ArH), 7.20-7.32 (6H, m, ArH), 7.42 (1H, d, J 7.7Hz, ArH),  
 5 and 10.93 (1H, br s, NH);  $m/z$  ( $CI^+$ ,  $NH_3$ ) 306 ( $M+1$ ) $^+$ .

The following were prepared in an analogous manner:

#### EXAMPLE 4

##### 10      2-(4-Phenylpiperazin-1-ylmethyl)indole

M.p. 264°C (toluene); (Found: C, 78.68; H, 7.35; N, 14.33.  $C_{19}H_{21}N_3$  requires C, 78.32; H, 7.26; N, 14.42%);  $\delta_H$  (DMSO- $d_6$ ) 2.55 (4H, t, J 4.9Hz, 2 x piperazinyl  $CH_2$ ), 3.14 (4H, t, J 4.9Hz, 2 x piperazinyl  $CH_2$ ), 3.66 (2H, s,  $CH_2$ -N), 6.29 (1H, s, 3-H), 6.75 (1H, t, J 7.3Hz, ArH), 6.89-6.95 (3H, m, ArH), 7.00 (1H, m, ArH), 7.16-7.24 (2H, m, ArH), 7.32 (1H, d, J 7.2Hz), 7.44 (1H, d, J 7.2Hz, ArH), and 11.00 (1H, br s, NH);  $m/z$  ( $CI^+$ ,  $NH_3$ ) 292 ( $M+1$ ) $^+$ .  
 15

#### EXAMPLE 5

##### 20      2-(4-[4-Chlorophenyl]piperazin-1-ylmethyl)indole hydrogen oxalate

M.p. 220°C (dec) (MeOH/ $H_2O$ ); (Found: C, 60.25; H, 5.23; N, 9.67.  $C_{19}H_{20}N_3Cl \cdot (CO_2H)_2$  requires C, 60.65; H, 5.33; N, 10.10%);  $\delta_H$  (DMSO- $d_6$ ) 2.77 (4H, m, 2 x piperazinyl  $CH_2$ ), 3.22 (4H, m, 2 x piperazinyl  $CH_2$ ), 3.91 (2H, s,  $NCH_2$ ), 6.40 (1H, s, 3-H), 6.93-6.99 (3H, m, ArH), 7.06 (1H, t, J 7.3Hz, ArH), 7.23 (2H, m, ArH), 7.35 (1H, d, J 7.6Hz, 4-H), 7.49 (1H, d, J 7.5Hz, 7-H), and 11.13 (1H, br s, NH);  $m/z$  ( $CI^+$ ,  $NH_3$ ) 326 ( $M+1$ ) $^+$ .  
 25

- 26 -

EXAMPLE 62-(4-Phenyl-1,2,3,6-tetrahydropyridin-1-ylmethyl)indole

M.p. 128-130°C (EtOAc); (Found: C, 82.14; H, 6.89; N, 9.46.  
 $C_{20}H_{20}N_2 \cdot 0.25H_2O$  requires C, 82.02; H, 7.06; N, 9.56%);  $\delta_H$   
 5 (DMSO- $d_6$ ) 2.48-2.50 (2H, m,  $NCH_2CH_2$ ), 2.69 (2H, t, J 5.6Hz,  
 $NCH_2CH_2$ ), 3.11 (2H, d, J 3.0Hz,  $NCH_2-CH=$ ), 3.73 (2H, s,  
 $NCH_2Ar$ ), 6.15 (1H, m,  $NCH_2-CH=$ ), 6.30 (1H, s, 3-H), 6.93 (1H,  
 m, ArH), 6.99-7.04 (1H, m, ArH), 7.23 (1H, t, J 7.2Hz, ArH),  
 7.29-7.34 (3H, m, ArH), 7.40-7.45 (3H, m, ArH), and 11.03 (1H,  
 10 br s, NH); m/z ( $CI^+$ ,  $NH_3$ ) 289 (M+1) $^+$ .

EXAMPLE 7

15 2-(4-[4-Methoxyphenyl]piperazin-1-ylmethyl)indole  
hydrogen oxalate

M.p. 218-220°C (dec) (MeOH); (Found: C, 64.55; H, 6.10;  
 N, 9.95.  $C_{20}H_{23}N_3O \cdot (CO_2H)_2$  requires C, 64.22; H, 6.12; N,  
 10.21%);  $\delta_H$  (DMSO- $d_6$ ) 2.89 (4H, br s, 2 x piperaziny CH<sub>2</sub>); 3.14  
 20 (4H, br s, 2 x piperaziny CH<sub>2</sub>), 3.67 (3H, s, OCH<sub>3</sub>), 4.05 (2H, s,  
 $NCH_2Ar$ ), 6.46 (1H, s, 3-H), 6.81 (2H, d, J 9.1Hz, ArH), 6.90 (2H,  
 d, J 9.1Hz, ArH), 6.99 (1H, t, J 7Hz, indole-H), 7.08 (1H, t, J  
 7Hz, indole-H), 7.37 (1H, d, J 8Hz, 4-H), 7.51 (1H, d, J 8Hz,  
 7-H), 11.25 (1H, br s, NH); m/z ( $CI^+$ ,  $NH_3$ ) 322 (M+1) $^+$ .

25

EXAMPLE 8

2-(4-[2-Phenylethyl]piperidin-1-ylmethyl)indole hydrogen  
oxalate

M.p. 230-232°C (dec) (MeOH); (Found: C, 69.98; H, 6.87; N,  
 30 6.81.  $C_{22}H_{26}N_2 \cdot (CO_2H)_2 \cdot 0.25H_2O$  requires C, 69.80; H, 6.96; N,

- 27 -

6.78%);  $\delta_{\text{H}}$  (DMSO- $d_6$ ) 1.38 (3H, m, aliphatic H), 1.51 (2H, m, aliphatic H), 1.86 (2H, m, aliphatic H), 2.55 (2H, m, aliphatic H), 2.67 (2H, m, aliphatic H), 3.22 (2H, m, aliphatic H), 4.19 (2H, s,  $\text{NCH}_2\text{Ar}$ ), 6.52 (1H, s, 3-H), 6.98-7.02 (1H, m, ArH), 7.09-7.19 (4H, m, ArH), 7.24-7.28 (2H, m, ArH), 7.40 (1H, d, J 9Hz, 4-H), 7.53 (1H, d, J 8Hz, 7-H), and 11.28 (1H, br s, NH);  $m/z$  ( $\text{CI}^+$ ,  $\text{NH}_3$ ) 319 ( $\text{M}+1$ ) $^+$ .

EXAMPLE 910      2-(4-[Benzyloxy]piperidin-1-ylmethyl)indole

M.p. 103-105°C (MeOH); (Found: C, 78.33; H, 7.48; N, 8.56.  $\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}$  requires C, 78.71; H, 7.55; N, 8.74%);  $\delta_{\text{H}}$  (DMSO- $d_6$ ) 1.54 (2H, m, piperidiny  $\text{CH}_2$ ), 1.85 (2H, m, piperidiny  $\text{CH}_2$ ), 2.12 (2H, t, J 9Hz, piperidiny  $\text{CH}_2$ ), 2.71 (2H, m, piperidiny  $\text{CH}_2$ ), 3.39 (1H, m, piperidiny CH), 3.57 (2H, s,  $\text{NCH}_2\text{Ar}$ ), 4.48 (2H, s,  $\text{OCH}_2\text{Ph}$ ), 6.24 (1H, s, 3-H), 6.92 (1H, m, ArH), 7.00 (1H, m, ArH), 7.23-7.35 (6H, m, ArH), 7.42 (1H, d, J 7.7Hz, 7-H), and 10.94 (1H, br s, NH);  $m/z$  ( $\text{CI}^+$ ,  $\text{NH}_3$ ) 321 ( $\text{M}+1$ ) $^+$ .

EXAMPLE 10

2-(4-[2-Phenylethyl]-1,2,3,6-tetrahydropyridin-1-ylmethyl)  
indole hydrogen oxalate hemihydrate

M.p. 226-228°C (dec) (MeOH); (Found: C, 69.59; H, 6.23; N, 6.71.  $\text{C}_{22}\text{H}_{24}\text{N}_2 \cdot (\text{CO}_2\text{H})_2 \cdot 0.5\text{H}_2\text{O}$  requires C, 69.38; H, 6.55; N, 6.74%);  $\delta_{\text{H}}$  (DMSO- $d_6$ ) 2.27 (4H, m, 2 x  $\text{CH}_2$ ), 2.69 (2H, m,  $\text{CH}_2$ ), 3.04 (2H, br s,  $\text{CH}_2$ ), 3.27 (2H, br s,  $\text{CH}_2$ ), 4.20 (2H, s,  $\text{NCH}_2\text{Ar}$ ), 5.41 (1H, s,  $\text{NCH}_2\text{-CH=}$ ), 6.51 (1H, s, 3-H), 7.00 (1H, t, J 7.3Hz, ArH), 7.08-7.37 (6H, m, ArH), 7.39 (1H, d, J 8Hz, 4-H), 7.52 (1H, d, J 8Hz, 7-H), and 11.34 (1H, br s, NH);  $m/z$  ( $\text{CI}^+$ ,  $\text{NH}_3$ ) 317 ( $\text{M}+1$ ) $^+$ .

- 28 -

EXAMPLE 112-(4-Phenylpropyl-1,2,3,6-tetrahydropyridin-1-ylmethyl)  
indole hydrogen oxalate

5 M.p. 208-210°C (EtOH); (Found: C, 71.27; H, 6.84; N,  
6.56.  $C_{23}H_{26}N_2 \cdot (CO_2H)_2$  requires C, 71.41; H, 6.71; N, 6.66%);  
 $\delta_H$  (DMSO- $d_6$ ) 1.67 (2H, m,  $PhCH_2CH_2CH_2$ ), 2.00 (2H, t, J 7.2Hz,  
10  $CH_2$ ), 2.22 (2H, br s, tetrahydropyridinyl  $CH_2$ ), 2.54 (2H, t, J  
7.6Hz,  $CH_2$ ), 3.06 (2H, br s, tetrahydropyridinyl  $CH_2$ ), 3.41 (2H,  
br s, tetrahydropyridinyl  $CH_2$ ), 4.24 (2H, s,  $NCH_2$ ), 5.39 (1H, s,  
tetrahydropyridinyl CH), 6.53 (1H, s, 3-H), 7.00 (1H, t, J 7.0Hz,  
ArH), 7.09-7.19 (4H, m, ArH), 7.26 (2H, t, J 7.3Hz, ArH), 7.39  
(1H, d, J 8.0Hz, 4-H), 7.53 (1H, d, J 7.8Hz, 7-H), and 11.38 (1H,  
15 br s, NH); m/z ( $CI^+$ ,  $NH_3$ ) 331 ( $M+1$ )<sup>+</sup>.

EXAMPLE 122-([4-(3'-Phenylpropylidene)piperidin-1-yl]methyl)indole  
hydrogen oxalate

20 M.p. 172-174°C (EtOH/ $H_2O$ ); (Found: C, 71.67; H, 6.68; N,  
6.74.  $C_{23}H_{26}N_2 \cdot (CO_2H)_2$  requires C, 71.41; H, 6.71; N, 6.66%);  
 $\delta_H$  (DMSO- $d_6$ ) 2.22-2.26 (6H, m, aliphatic H), 2.56-2.60 (4H, m,  
aliphatic H), 2.79 (2H, br s,  $CH_2$ ), 4.08 (2H, s,  $NCH_2$ ), 5.24 (1H, t,  
25 J 7.1Hz,  $PhCH_2CH_2CH$ ), 6.45 (1H, s, 3-H), 6.98-7.20 (7H, m,  
ArH), 7.39 (1H, d, J 8.1Hz, 4-H), 7.52 (1H, d, J 7.8Hz, 7-H), and  
11.29 (1H, br s, NH); m/z ( $CI^+$ ,  $NH_3$ ) 331 ( $M+1$ )<sup>+</sup>.

- 29 -

EXAMPLE 132-(1,2,3,4-Tetrahydroisoquinolin-2-ylmethyl)indole

5 M.p. 129-131°C (Me<sub>2</sub>CO/H<sub>2</sub>O); (Found: C, 81.91; H, 6.87; N, 10.57. C<sub>18</sub>H<sub>18</sub>N<sub>2</sub> · 0.1(H<sub>2</sub>O) requires C, 81.84; H, 6.94; N, 10.61%); δ<sub>H</sub> (CDCl<sub>3</sub>) 2.84 (2H, t, J 5.8Hz, PhCH<sub>2</sub>CH<sub>2</sub>N), 2.94 (2H, t, J 5.8Hz, PhCH<sub>2</sub>CH<sub>2</sub>N), 3.69 (2H, s, PhCH<sub>2</sub>N), 3.89 (2H, s, ArCH<sub>2</sub>N), 6.41 (1H, t, J 0.9Hz, 3-H), 6.96 (1H, d, J 7.0Hz, ArH),  
10 7.06-7.17 (5H, m, ArH), 7.32 (1H, d, J 8.0Hz, 4-H), 7.57 (1H, d, J 7.6Hz, 7-H), and 8.70 (1H, br s, NH); m/z (CI<sup>+</sup>, NH<sub>3</sub>) 263 (M+1)<sup>+</sup>.


EXAMPLE 142-(1,3-Dihydroisoindol-2-ylmethyl)indole

M.p. 176-178°C (dec) (MeOH/H<sub>2</sub>O); (Found: C, 82.58; H, 6.42; N, 11.13. C<sub>17</sub>H<sub>16</sub>N<sub>2</sub> requires C, 82.22; H, 6.49; N, 11.28%); δ<sub>H</sub> (CDCl<sub>3</sub>) 3.97 (4H, s, 2 x dihydroisoindolyl CH<sub>2</sub>), 4.07 (2H, s, NCH<sub>2</sub>Ar), 6.40 (1H, t, J 0.9Hz, 3-H), 7.06-7.22 (7H, m, ArH), 7.28 (1H, dd, J 7.9, 0.8Hz, 4-H), 7.57 (1H, m, 7-H), and 8.64 (1H, br s, NH); m/z (CI<sup>+</sup>, NH<sub>3</sub>) 249 (M+1)<sup>+</sup>.

- 30 -

**CLAIMS:**

1. The use of a compound of formula I, or a  
pharmaceutically acceptable salt thereof or a prodrug  
5 thereof:




( I )

wherein


E represents  $-\text{CH}_2-$  or  $-\text{CH}_2\text{CH}_2-$ ;R represents hydrogen or  $\text{C}_{1-6}$  alkyl;

Q represents a moiety of formula Qa, Qb, Qc or


20 Qd:




( Q a )



( Q b )



( Q c )



( Q d )



- 31 -

in which the broken line represents an optional chemical bond;

5  $R^1$  represents hydrogen, or an optionally substituted  $C_{1-6}$  alkyl,  $C_{1-6}$  alkoxy,  $C_{2-6}$  alkenyl,  $C_{2-6}$  alkynyl, aryl, aryl( $C_{1-6}$ )alkyl, aryl( $C_{1-6}$ )alkoxy, aryl( $C_{2-6}$ )alkenyl, aryl( $C_{2-6}$ )alkynyl,  $C_{3-7}$  heterocycloalkyl( $C_{1-6}$ )alkyl, heteroaryl, heteroaryl( $C_{1-6}$ )alkyl, heteroaryl( $C_{2-6}$ )alkenyl or heteroaryl( $C_{2-6}$ )alkynyl group;

10  $R^2$  represents an optionally substituted  $C_{1-6}$  alkyl,  $C_{1-6}$  alkoxy,  $C_{2-6}$  alkenyl,  $C_{2-6}$  alkynyl, aryl, aryl( $C_{1-6}$ )alkyl, aryl( $C_{1-6}$ )alkoxy, aryl( $C_{2-6}$ )alkenyl, aryl( $C_{2-6}$ )alkynyl,  $C_{3-7}$  heterocycloalkyl( $C_{1-6}$ )alkyl, heteroaryl, heteroaryl( $C_{1-6}$ )alkyl, heteroaryl( $C_{2-6}$ )alkenyl or heteroaryl( $C_{2-6}$ )alkynyl group;

15  $R^3$ ,  $R^4$  and  $R^5$  independently represent hydrogen, hydrocarbon, a heterocyclic group, halogen, cyano, trifluoromethyl, nitro,  $-OR^a$ ,  $-SR^a$ ,  $-SOR^a$ ,  $-SO_2R^a$ ,  $-SO_2NR^aR^b$ ,  $-NR^aR^b$ ,  $-NR^aCOR^b$ ,  $-NR^aCO_2R^b$ ,  $-COR^a$ ,  $-CO_2R^a$  or  $-CONR^aR^b$ ;


20  $Z$  represents  $-CH_2-$  or  $-CH_2CH_2-$ ;

$R^6$  represents hydrogen,  $C_{1-6}$  alkyl,  $C_{1-6}$  alkoxy, aryl, aryl( $C_{1-6}$ )alkyl or halogen; and

25  $R^a$  and  $R^b$  independently represent hydrogen, hydrocarbon or a heterocyclic group; for the manufacture of a medicament for the treatment and/or prevention of psychotic disorders.

30 2. The use as claimed in claim 1 of a compound represented by formula IIA, and pharmaceutically acceptable salts thereof and prodrugs thereof:

- 32 -



(IIA)

wherein

E is as defined in claim 1;

n is zero, 1, 2 or 3;

W is CH or nitrogen; and

15  $R^{13}$  and  $R^{17}$  independently represent hydrogen, halogen, cyano, nitro, trifluoromethyl, amino,  $C_{1-6}$  alkylamino, di( $C_{1-6}$ )alkylamino,  $C_{1-6}$  alkyl,  $C_{1-6}$  alkoxy, aryl( $C_{1-6}$ )alkoxy or  $C_{2-6}$  alkylcarbonyl.

20 3. The use as claimed in claim 1 of a compound selected from:

2-(4-phenylpiperazin-1-ylmethyl)-1H-indole;

2-[4-(4-chlorophenyl)piperazin-1-ylmethyl]-1H-indole;

25 and pharmaceutically acceptable salts thereof and prodrugs thereof.

4. A method for the treatment and/or prevention of psychotic disorders, which comprises administering to a patient in need of such treatment an effective amount of a compound of formula I as defined in claim 1, or a pharmaceutically acceptable salt thereof or a prodrug thereof.

30

5. The method as claimed in claim 4 wherein the compound administered is represented by formula IIA


35

- 33 -

as defined in claim 2, or a pharmaceutically acceptable salt thereof or a prodrug thereof.

6. The method as claimed in claim 4 wherein the compound administered is selected from:  
 2-(4-phenylpiperazin-1-ylmethyl)-1H-indole;  
 2-[4-(4-chlorophenyl)piperazin-1-ylmethyl]-1H-indole;  
 and pharmaceutically acceptable salts thereof and prodrugs thereof.

7. A compound of formula IIB, or a salt or prodrug thereof:




(IIB)

wherein

E is as defined in claim 1; and  
 n, R<sup>13</sup> and R<sup>17</sup> are as defined in claim 2.

8. A compound of formula IIC, or a salt or prodrug thereof:



(IIC)

- 34 -

wherein


m is 1, 2 or 3;

-X-Y- represents  $-\text{CH}_2\text{CH}-$  or  $-\text{CH}=\text{C}-$ ;

E is as defined in claim 1; and

5  $\text{R}^{13}$  and  $\text{R}^{17}$  are as defined in claim 2.

9. A compound of formula IID, or a salt or prodrug thereof:



(IID)

wherein


E is as defined in claim 1;

20 Z represents  $-\text{CH}_2-$  or  $-\text{CH}_2\text{CH}_2-$ ;

$\text{R}^{13}$  is as defined in claim 2; and

$\text{R}^{16}$  represents hydrogen,  $\text{C}_{1-6}$  alkyl,  $\text{C}_{1-6}$  alkoxy, aryl, aryl( $\text{C}_{1-6}$ )alkyl or halogen.

25 10. A compound of formula IIE, or a salt or prodrug thereof:



(IIE)

wherein

- 35 -

E is as defined in claim 1;  
R<sup>13</sup> and R<sup>17</sup> are as defined in claim 2; and  
m is 1, 2 or 3.

- 5                   11. A compound selected from:
- 2-(4-benzylpiperazin-1-ylmethyl)-1H-indole;  
2-[2-(4-benzylpiperazin-1-yl)ethyl]-1H-indole;  
2-[4-(4-methoxyphenyl)piperazin-1-ylmethyl]-1H-indole;  
10 2-(4-phenyl-1,2,3,6-tetrahydropyrid-1-ylmethyl)-1H-  
indole;  
2-[4-(2-phenylethyl)piperidin-1-ylmethyl]-1H-indole;  
2-(4-benzylloxypiperidin-1-ylmethyl)-1H-indole;  
2-[4-(2-phenylethyl)-1,2,3,6-tetrahydropyrid-1-ylmethyl]-  
1H-indole;  
15 2-[4-(3-phenylpropyl)-1,2,3,6-tetrahydropyrid-1-  
ylmethyl]-1H-indole;  
2-[4-(3-phenylpropylidene)piperidin-1-ylmethyl]-1H-  
indole;  
2-(1,2,3,4-tetrahydroisoquinolin-2-ylmethyl)-1H-indole;  
20 2-(1,2-dihydroisoindol-2-ylmethyl)-1H-indole;  
and salts and prodrugs thereof.

12. A pharmaceutical composition comprising a  
compound as claimed in any one of claims 7 to 11 in  
25 association with a pharmaceutically acceptable carrier.

13. A compound as claimed in any one of claims  
7 to 11 for use in therapy.

- 30 14. The use of a compound as claimed in any  
one of claims 7 to 11 for the manufacture of a medicament  
for the treatment and/or prevention of psychotic  
disorders.

- 35 15. A method for the treatment and/or  
prevention of psychotic disorders, which comprises

- 36 -

administering to a patient in need of such treatment an effective amount of a compound as claimed in any one of claims 7 to 11.

# INTERNATIONAL SEARCH REPORT

Inter      nal Application No  
PCT/GB 94/00760

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 5    C07D209/14    A61K31/40    C07D401/06

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 5    C07D    A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL ABSTRACTS, vol. 66, no. 7, 13 February 1967, Columbus, Ohio, US; abstract no. 28694r, K.T. POTTS ET AL. 'Synthetic experiments related to the indole alkaloids. V. ...' see abstract * RN 102892-13-2 * & J. HETEROCYCL. CHEM. 3(4), 395-401 ----	9
A	EP,A,0 411 631 (MERREL DOW PHARMACEUTICALS INC.) 6 February 1991 see claims ----	1
A	US,A,3 188 313 (STERLING DRUG INC.) 8 June 1965 cited in the application * column 15, line 3-15; claim 21 * -----	1

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*&\* document member of the same patent family

Date of the actual completion of the international search

23 June 1994

Date of mailing of the international search report

13. 07. 94

Name and mailing address of the ISA

European Patent Office, P.O. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. ( + 31-70 ) 340-2040, Tlx. 31 651 epo nl,  
Fax ( + 31-70 ) 340-3016

Authorized officer

Van Bijlen, H

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/GB94/00760

**Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)**

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
Although claims 4 - 6 and 15 are directed to a method of treatment of (diagnostic method practised on) the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

**Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)**

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.



# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PC1/GB 94/00760

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0411631	06-02-91	AU-B- 625238	02-07-92
		AU-A- 6012390	07-02-91
		CA-A- 2022451	04-02-91
		DE-D- 69008069	19-05-94
		JP-A- 3066683	22-03-91
		US-A- 5064840	12-11-91
-----			
US-A-3188313		DE-A- 1445151	09-01-69
		GB-A- 944443	
-----			